

# Birth weight pattern and factors affecting birth weight in urban areas of South-Western Ethiopia

Seleshi T/Mariam<sup>1</sup>, Ahmed Ali<sup>2</sup>

**Abstract:** A community-based cross-sectional study of birth-weight was carried out in 30 villages of Jimma Zone, to determine the pattern of birth weight, its relationship with ante-natal care, and the influencing factors in birth-weight. A total of 537 mother-newborn pairs were included in the study revealing an overall low birth weight rate of 10.6%.

About 72% of the mothers were booked for ante-natal care during the current pregnancy, and 332 (86%) of the deliveries were attended by trained personnel, including trained traditional birth attendants. Mean birth-weight for both sexes was 3,202 grams; and there was no statistically significant difference between the sexes ( $P>0.05$ ). Higher mean birth weight was observed for newborns whose mothers attended ante-natal clinic and whose births were attended by trained personnel than those mothers without ante-natal care and unattended by trained personnel ( $P<0.05$ ). Low birth weight was associated with being single, poor, without ante-natal care and not attended by trained personnel during delivery. Efforts to increase the utilization of ante-natal care services is recommended to minimize the risk of low birth-weight as it will help the timely correction of factors predicting low birth weight. [*Ethiop. J. Health Dev.* 1998;12(1):33-37]

## Introduction

Birth-weight is governed by two major factors: duration of gestation and intrauterine growth rate. Antenatal care (ANC) may benefit both factors through early identification and proper management of risk factors, and pregnancy complications. It is very difficult to reach at firm conclusions about the possible benefit of early or frequent antenatal visits on either intrauterine growth or gestational duration (1). There are a number of components of ANC interventions that have been found to be effective, investigation, detection, treatment and referral of anaemia, hypertension and infections being some in the list. Therefore, foetal weight gain, birth-weight, and neonatal health have been shown to be significantly affected by the mother's use of ANC. Nevertheless, only a small number of the population in developing countries receive ANC (2). In Ethiopia coverage with ANC ranges from around 10% in rural areas to 60% in some urban areas. Data on distribution of birth weight are scanty. The few available are from deliveries conducted by health institutions. Of all deliveries in Ethiopia, more than 85% are attended by lay personnel and take place outside health units. In addition to having problems of coverage and representativeness, data from health units lack validity and reliability (3-6). Although they are inconsistent in defining low birth-weight (LBW), the existing institution-based data indicate a high magnitude of LBW in Ethiopia (7-11).

WHO estimates the incidence of LBW for Ethiopia as 10-20% (3). A similar estimate made by UNICEF indicates the percentage of LBW in Ethiopia to be 13% with the average birth-weight being 10% lower than those born in some developed countries (6).

The objectives of this study were to assess the magnitude of LBW, its relationship with

---

<sup>1</sup>From the Department of Community Health, Jimma Institute of Health Sciences, P.O. Box 378, Jimma, Ethiopia and <sup>2</sup>Department of Community Health, Faculty of Medicine Addis Ababa University.

prenatal care and to examine factors governing birth-weight among delivered newborns in Jimma Zone. It is hoped that findings of this study will provide better and representative data on the pattern of birth-weight and factors affecting it.

## Methods

**Study area:** This cross-sectional community-based study was undertaken in 30 villages of Jimma

administrative zone in South Western Ethiopia. The villages were selected using convenience sampling techniques. Two health centers, three clinics, and one hospital were located within 10 and 50 kms. radius from these villages, respectively. All the health facilities are involved in providing delivery and antenatal clinic (ANC) services.

*Study population:* All delivered newborns from 1st October to 31st January, 1993 were included in the study.

*Data collection and analysis:* Expectant mothers were registered, and visited on daily basis throughout their 3rd trimester by TTBAAs and enumerators of their respective villages. The TTBAAs and the enumerators met twice daily (morning and afternoon) in their respective Kebele offices and exchanged information on women who have delivered. A pre-tested structured questionnaire consisting of socio-economic, obstetric, and MCH service utilization pattern of the mother was completed by trained female enumerators within 24 hours after birth. The enumerators were provided with a salter spring balance with a capacity to weigh up to 10,000 grams and a piece of cloth weighing less than 5 grams to be used as a weighing pant. The accuracy of the balances was checked regularly using a standard weight.

The National Center for Health Statistics (NCHS) reference was used to define LBW i.e <2500 gm or less (12). Women who had had one or more visits to an ANC clinic in the current pregnancy were considered as users of ANC. Still births and multiple deliveries were excluded for analysis of this study. Data were entered and analyzed using SPSS/PC statistical package. Descriptive statistics as well as analytic methods were employed during analysis and presentation of the data.

## Results

A total of 543 newborns were reported during the study period. Of these, six (two stillbirths and two sets of twin deliveries) were excluded while the remaining 537 mother- newborn pairs were included in this study. Of the 537 singleton live births reported, 393(73.1%), 123(22.9%) and 21(3.9%) were from Jimma, Agaro, Yebu and Serbo towns, respectively.

Table 1: **Distribution of study mothers by socio-economic characteristics, Jimma, 1994.**

| Characteristics      | N   | %     |
|----------------------|-----|-------|
| Marital status       |     |       |
| Married              | 484 | 90.1  |
| Unmarried            | 53  | 9.9   |
| Total                | 537 | 100.0 |
| Educational status   |     |       |
| Can't read and write | 153 | 28.5  |
| 1-8 grade            | 170 | 31.7  |
| Grade 9 & above      | 214 | 39.8  |
| Total                | 537 | 100.0 |
| Occupational status  |     |       |
| House wife           | 442 | 82.5  |
| Others               | 94  | 17.5  |
| Total*               | 536 | 100.0 |
| Family Income        |     |       |
| No. Income           | 50  | 9.3   |
| ≤100 Birr            | 181 | 33.7  |
| 101-200 Birr         | 137 | 25.5  |
| 201-300 Birr         | 67  | 12.5  |
| 301- 400 Birr        | 28  | 5.2   |
| 400*                 | 74  | 13.8  |
| Total                | 537 | 100.0 |

\* Analysis is based on non-missing data.

The mean age of the mothers was 25 years with a SD of 5.9. Among the study mothers 24.7% were primiparous, 27.7% grand multiparous (≥5) and 47.6% were with 2-4 previous pregnancies. History of prior abortion and stillbirths were reported in about 11% and 2% of the study mothers, respectively. The majority of the mothers were married (90.1%), able to read and write (72.9%),

and house wives (82.5%) (Table 1). Table 2 shows the distribution of newborns by sex and birth-

Table 2: **Distribution of new borns by sex and category of birth weight Jimma, 1994.**

| Category of Birth Weight | Male |      | Female |      | Total |       |
|--------------------------|------|------|--------|------|-------|-------|
|                          | N    | %    | N      | %    | N     | %     |
| Normal birth weight      | 273  | 57.9 | 207    | 43.1 | 480   | 89.4  |
| Low birth weight         | 28   | 49.1 | 29     | 51.9 | 57    | 10.6  |
| Total                    | 301  | 56.1 | 236    | 43.9 | 537   | 100.0 |

$\chi^2 = 1.24$

$P = 0.2655$

LBW < = 2500 gms;

NBW > 2500 gms

weight category. As indicated in the table the overall LBW rate was 10.6% with a male to female ratio of 1.3:1. Comparison of normal and low birth-weights by socio-demographic and health service use factors is presented in Table 3. LBW babies were found mostly among single and low family income mothers than others ( $P < 0.05$ ). A higher proportion of mothers with LBW babies were not booked for ANC and their deliveries were unattended by trained personnel than did women whose pregnancy ended in normal birth-weight (NBW) babies. The difference is statistically significant ( $P < 0.05$ ).

Table 4 summarizes the distribution of the study mothers' ANC status by attendants of the delivery. Trained personnel, including TTBA's attended 81.4% of the deliveries. There was a significant difference between attended and unattended deliveries with the status of antenatal care ( $P < 0.01$ ). Table 5 shows the influence of different socio-demographic variables on mean birth-weight. Newborns delivered to mothers who were married and have visited an ANC at least once during the current pregnancy had higher mean birth-weight than newborns whose mothers were single and did not have ANC follow-up ( $P < 0.05$ ). However mean birth-weight was not influenced by factors like literacy status of mothers and sex of the baby ( $P > 0.05$ ).

*Results of Regression analysis:* Those independent variables, having significant bi-variate association with birth-weight in Tables 3 and 5 were entered into a linear regression model in consecutive steps. Among these variables used in the linear regression model only ANC and delivery attendant were found to have a significant relationship with birth weight at  $p = 0.02$  and  $p = 0.001$ , respectively. The overall contribution of the variables when taken together is estimated to be .31047 (the linear regression explains a significant proportion of the total variation in birth-weight). However other factors which were not included might affect birth weight in this study.

## Discussion

Though strict comparisons may be difficult, the overall prevalence of LBW observed in this study is close to the reports from Gondar and Jimma (9,11). The overall mean birth weight in this study is slightly higher than the previous report from Jimma (11). Males in this study tended to have lower mean birth-weight than their female counterparts when compared to other studies in Jimma (10,11). The likely explanation for the higher LBW rate in those mothers who were not married and whose deliveries were not attended by trained personnel is that they are more likely to be of lower income group and may have less access to health care facilities. Furthermore, they might make less use of ANC and their nutritional status may also be sub-optimal as reported in other previous studies (13,14). As reported elsewhere (15,16), the more socio-economically disadvantaged the subpopulation, the poorer the perinatal outcome, and the higher the incidence of LBW. This conforms with the association of family income level and LBW in our study. On the contrary, other studies have reported that income did not differ

**Table 3: Comparison of normal and low birth-weight babies by socio-demographic, and health service use factors, Jimma, 1994.**

| Characteristics      | Birth weight category |      |     |      |                |         |
|----------------------|-----------------------|------|-----|------|----------------|---------|
|                      | LBW                   |      | NBW |      | X <sup>2</sup> | P-value |
|                      | N                     | %    | N   | %    |                |         |
| ANC                  |                       |      |     |      |                |         |
| Yes                  | 23                    | 40.4 | 128 | 26.7 | 4.96           |         |
| No                   | 34                    | 59.6 | 352 | 73.3 |                | P=0.025 |
| Birth attendant      |                       |      |     |      |                |         |
| Trained              | 38                    | 66.7 | 397 | 82.7 | 7.51           |         |
| Untrained            | 19                    | 33   | 83  | 17.3 |                | P=0.006 |
| Place of delivery    |                       |      |     |      |                |         |
| Health Unit          | 20                    | 35.1 | 231 | 48.1 | 2.97           |         |
| Home                 | 37                    | 64.9 | 249 | 51.9 |                | P=0.084 |
| Marital status       |                       |      |     |      |                |         |
| Married              | 48                    | 84.2 | 453 | 94.4 | 6.86           |         |
| Unmarried            | 9                     | 15.8 | 27  | 5.6  |                | p=0.009 |
| Literacy status      |                       |      |     |      |                |         |
| Grade 1 & above      | 19                    | 33.3 | 353 | 23.7 | 0.94           |         |
| Can't read and write | 38                    | 66.7 | 126 | 26.3 |                | p=0.332 |
| Family Income        |                       |      |     |      |                |         |
| No Income            | 10                    | 17.5 | 40  | 8.3  |                |         |
| <100                 | 25                    | 43.9 | 156 | 32.5 |                |         |
| 101-200              | 12                    | 21.1 | 126 | 26.3 | 11.20          |         |
| 201-300              | 3                     | 5.3  | 63  | 13.1 |                |         |
| 301-400              | 2                     | 3.5  | 26  | 5.4  |                | p=0.024 |
| 400+                 | 5                     | 8.8  | 69  | 14.4 |                |         |
| Sex of Baby          |                       |      |     |      |                |         |
| Male                 | 28                    | 49.1 | 273 | 56.9 |                |         |
| Female               | 29                    | 50.9 | 207 | 43.1 |                | p=0.33  |

LBW <= 2500 gms; NBW > 2500 gms

significantly between LBW and NBW mothers (1). The provision of ANC is a means for better utilization of trained personnel and health units for delivery services as it was found in our study that mothers who had ANC are more likely to deliver in health units or by trained personnel than mothers without ANC. Among important factors included in this study, age, parity, religion and ethnicity of the mother were not found to be associated with birth-weight of an infant. The factors mentioned were among factors predicting birth-weight in other studies (1,6,18,19,20). The observed non-significant difference might be due to the relatively better maternal level of education in our study where mothers may be more likely to comply with medical advice during pregnancy or to make better use of health care services.

**Table 4: Distribution of mothers ANC status by attendant of delivery, Jimma, 1994.**

| Attendant of delivery | Antenatal Clinic |      |     |      |       |      |
|-----------------------|------------------|------|-----|------|-------|------|
|                       | Yes              |      | No  |      | Total |      |
|                       | N                | %    | N   | %    | N     | %    |
| Trained               | 332              | 86.0 | 115 | 69.5 | 437   | 81.4 |
| Untrained             | 54               | 14.0 | 46  | 0.5  | 00    | 8.6  |
| Total                 | 386              | 71.9 | 151 | 28.1 | 537   | 00.0 |

$\chi^2=16.17$

P=0.0005

Although gestational age and prematurity were not determined, bias as a possible explanation of the finding of the study was minimized by providing training to enumerators, and through frequent supervision, to avoid measurement bias during data collection. Moreover data quality was controlled by five field supervisors and the principal investigator regularly.

Table 5: Mean birth-weight and factors affecting mean birth-weight, Jimma, 1994.

| Variables          | Mean birth weight | t-value | p-value  |
|--------------------|-------------------|---------|----------|
| ANC                |                   |         |          |
| Yes                | 3179.2            |         |          |
| No                 | 3071.9            |         |          |
| Birth attendant    |                   | 2.32    | p=0.0104 |
| Trained            | 3179.2            |         |          |
| Untrained          | 3071.2            |         |          |
| Place of delivery  |                   | 3.25    | p=0.0006 |
| Health unit        | 3191.2            |         |          |
| Home               | 3111.2            |         |          |
| Marital status     |                   | 2.12    | p=0.017  |
| Married            | 3162.1            |         |          |
| Not married        | 2961.1            |         |          |
| Literacy status    |                   | 3.48    | p=0.0002 |
| Grade 1 & above    | 3154.5            |         |          |
| Can't read & write | 3130.3            |         |          |
| Sex of baby        |                   | 0.54    | p=0.295  |
| Male               | 3169.1            |         |          |
| Female             | 3122.5            |         |          |

## Conclusion and Recommendation

The findings of this study have demonstrated that LBW is higher and mean birth-weight is lower among mothers without ANC. Based on this, we recommend that efforts to encourage mothers to utilize ANC services to minimize the risk of LBW is mandatory. Further study to measure the quality or quantity of ante-natal care is also recommended.

## References

1. Kramer MS. Determinants of Low Birth Weight. Methodologic Assessment and Meta-Analysis. Bull. WHO 1987;65:663-737.
2. Mesham D. Towards the development of safe motherhood programme guidelines. Report of a workshop organized by the World Bank and the mother care project. Family care International, 1992.
3. WHO. The Incidence of Low Birth Weight. A critical Review of Available Information. World Health Stat. Quart. 1980;22:197-224.
4. UNICEF. The state of the world children. Oxford University Press. New York 1991.
5. MOH. Health Service Directory. Planning & Programming Bureau. 1981(1989).
6. UNICEF. Children and Women in Ethiopia. Situational Analysis Addis Ababa. 1989.
7. Abate CG. Changes in Birth Weight Distribution from 1973 to 1982 in Addis Ababa. Bull WHO 1986;64:711-714.
8. Zein A. Zein. Birth weight of Hospital Delivered Neonates in Gondar North Western Ethiopia. Ethiop Med J 1985;23:59-64.
9. Zein A. Zein, Worku A. The distribution of LBW in Gondar Adm. Region, North Western Ethiopia. The Ethiopian Journal of Health Dev. 1991;5:71-74.
10. Kebede A. Health Consequences of Low Birth Weight in Ethiopia MPH Thesis to AAU 1989. (Unpublished)
11. Shiferaw T. Some Factors Associated with Birth weight in Jimma South Western Ethiopia. Ethiop Med J. 1990;28:83-189.
12. Kessele S, Villar J, Berendes HW, Nugent RP. The changing pattern of LBW in USA. J AM

Med Ass. 1984;251:1978-1982.

13. Jeffrey B, Gould Leroy S. Socio Economic Status, Low Birth Weight and Race. *Paediatrics* 1988;82:896-905.
14. Naeye RL, Taferi N, Gokezie A. Effects of Maternal under nutrition and Heavy physical Work During Pregnancy of Birth Weight. *Br. J. Ob Gyn.* 1980;87:222.
15. Barnes LB, Linda SA, Barry MP. Womens Physical Activity Pregnancy Outcome. A longitudinal Analysis from Philippines *Int J Epidemiol.* 1991;20:162-172.
16. Osafu, Wlkaija. Birth Weight of Babies In Benin: Bendal State of Nigeria. *J Trop Paediatrics.* 1985;31:139-143.
17. WHO. Development of Indicators for Monitoring Progress Towards HFA/2000. HFA series 1981, No. 4 Geneva.
18. Chen R, Waxy Lusky A, Toppleber, Bard V. Acrriterion for standard Definition of Low Birth Weight *Int J of Epidemiol.* 1991;20:180-186.
19. Padburg JF, Roderman B, Oddie TH, Hobel CJ, Fisher DA. Maternal Risk Factors In Infants with very Low Birth Weight. *Obs and Gyn.* 1982;60:612-616.
20. Kwang SL, Ferugson RM, Corpuz M, Gartner LM. Maternal age and the Incidence of Low birth weigh at Term. *Am J Obs Gyn* 1988;158:84-89.